

Temolina[®]

Linagliptin

COMPOSITION:

Temolina[®] 5 Tablet: Each film coated tablet contains Linagliptin INN 5 mg.

PHARMACOLOGY:

Linagliptin is a dipeptidyl peptidase-4 (DPP-4) inhibitor, which exerts its action by slowing the inactivation of incretin hormones. Incretin hormones, including glucagon-like peptide-I (GLP-I) and glucose-dependent Insulinotropic polypeptide (GIP), are released by the intestine throughout the day and levels are increased in response to a meal. These hormones are rapidly inactivated by the enzyme, DPP-4. The incretins are part of an endogenous system involved in the physiologic regulation of glucose homeostasis. When blood glucose concentrations are normal or elevated, GLP-I and GIP increase Insulin synthesis and release from pancreatic beta cells by intracellular signaling pathways involving cyclic AMP. GLP-I also lowers glucagon secretion from pancreatic alpha cells, leading to reduced hepatic glucose production.

INDICATION:

Linagliptin is indicated in the treatment of type-2 diabetes mellitus to improve glycemic control in adults:

as monotherapy :

- * In patients inadequately controlled by diet and exercise alone and for whom Metformin is inappropriate due to intolerance or contraindicated due to renal impairment.

as combination therapy :

- * In combination with Metformin when diet and exercise plus Metformin alone do not provide adequate glycaemic control.
- * In combination with a Sulphonylurea and Metformin when diet and exercise plus dual therapy with these medicinal products do not provide adequate glycemic control.
- * In combination with Insulin with or without Metformin, when this regimen alone, with diet and exercise, does not provide adequate glycemic control.

DOSE & ADMINISTRATION:

- * The recommended dose of Linagliptin is 5 mg once daily.
- * Linagliptin can be taken with or without food at any time of the day. If a dose is missed, it should be taken as soon as the patient remembers. A double dose should not be taken on the same day.
- * For patients with renal insufficiency no dosage adjustment is required.
- * Pharmacokinetic studies suggest that no dose adjustment is required for patients with hepatic impairment.
- * No dose adjustment is necessary based on age.
- * Clinical experience in patients >80 years of age is limited and caution should be exercised when treating this population.

CONTRA-INDICATION:

Linagliptin should not be prescribed for patients with a history of a hypersensitivity reaction to this drug, such as urticaria, angioedema, or bronchial hyper-reactivity.

WARNING AND PRECAUTION:

General - Linagliptin should not be used in patients with type - I diabetes or for the treatment of diabetic ketoacidosis.

Hypoglycemia - Linagliptin alone showed a comparable incidence of hypoglycemia to placebo. Caution is advised when Linagliptin is used in combination with a Sulphonylurea and/or Insulin. A dose reduction of the Sulphonylurea or Insulin, may be considered.

Pancreatitis - In post-marketing experience of Linagliptin there have been spontaneously reported adverse reactions of acute pancreatitis. Patients should be informed of the characteristic symptom of acute pancreatitis: persistent, severe abdominal pain. Resolution of pancreatitis has been observed after discontinuation of Linagliptin. If pancreatitis is suspected, Linagliptin should be discontinued.

SIDE EFFECTS:

The most common side effects of Linagliptin are stuffy or runny nose and sore throat. Hypoglycemia may occur when Linagliptin is combined with Insulin or Sulfonylurea and Metformin. Allergic reaction and muscle pain also may occur. Pancreatitis, angioedema (frequency rare) and urticaria (frequency rare) were identified as additional adverse reactions.

USE IN PREGNANCY & LACTATION:

Pregnancy: The use of Linagliptin has not been studied in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. As a precautionary measure, it is preferable to avoid the use of Linagliptin during pregnancy.

Nursing mothers: Available pharmacokinetic data in animals have shown excretion of Linagliptin metabolites in milk. A risk to the breast-feed child cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Linagliptin therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

USE IN CHILDREN & ADOLESCENTS:

The safety and efficacy of Linagliptin in children and adolescents has not yet been established. No data are available.

DRUG INTERACTION:

The efficacy of Linagliptin may be reduced when the drug is coadministered with a strong CYP3A4 or P-glycoprotein inducer (e.g., rifampin). Therefore, the use of alternative treatments is strongly recommended. Sulfonylureas should be used with caution during treatment with Linagliptin. The pharmacokinetic characteristics of Linagliptin were not altered by the concomitant administration of Simvastatin, Digoxin, Glyburide, Warfarin, Metformin or Pioglitazone.

OVERDOSE:

During controlled clinical trials in healthy subjects, single doses of up to 600 mg Linagliptin (equivalent to 120 times the recommended dose) were generally well tolerated. There is no experience with doses above 600 mg in humans.

Therapy in the event of an overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring and institute clinical measures if required.

STORAGE:

Store in a dry and cool place below 30° C temperature and keep away from light and moisture. Keep out of reach of children.

PACKING:

Temolina[®] 5 Tablet: Each box containing 2 x 15's Tablets in Alu-Alu blister pack.